


Brown Dog Tick

Control and prevention in domestic situations



Oct 2026

Chartered Institute of
Environmental Health



UK Health
Security
Agency

Brown Dog Tick

(Rhipicephalus sanguineus)

Control and prevention in domestic situations

In the UK, non-native ticks are imported on dogs and some of these result in subsequent infestation of owner's houses, kennels or other sheltered environments. The tick species associated with infestations is the brown dog tick. Unlike ticks commonly found in the UK, this species thrives indoors, and may require pest control intervention. Infestations present a potential health risk to animals and humans, due to the ability of this tick species to transmit pathogens that cause diseases.

Pet owners should be vigilant and inspect their animal(s) for tick activity, particularly dogs that have recently travelled abroad or been imported; areas of the animal prone to tick bites are - the ears, where the hind leg meets the body and the paws. However, a thorough check of the whole dog is required to ensure that all feeding ticks are found and removed with a tick remover tool/tweezers as soon as possible.

The host animal should be treated with an appropriate approved control product if ticks are present; pet owners should discuss treatment options with their veterinarian, letting them know they've been exposed to ticks from a different country. The owner of the animal or a veterinary surgeon should carry out this treatment.

Sometimes UK tick species can be found in dog bedding. However ticks found in furniture, on curtains, or climbing walls are likely to be the non-native species and require further investigation and control. Should tick infestation in a dwelling be suspected, pet owners should seek out expert identification via the UK Health Security Agency (see imported tick information via <https://www.gov.uk/guidance/tick-surveillance-scheme> or contact tick@ukhsa.gov.uk), so it is clear which species of tick is being dealt with. This accurate identification is important to determine what course of action may be needed regarding control.


Control of ticks in the home should be undertaken by a professional pest control operator (PCO). Treatment by a PCO would mainly involve the application of residual insecticides (acaricides) approved for tick control both indoors and around buildings. Such products can be sprayed directly on to ticks or to surfaces. Certain dessicant powders and Ultra Low Volume (ULV) cold fog products / techniques are also approved for tick control.

PCOs are trained to use these products and they would likely treat wall-floor-junctions, cracks & crevices where ticks may be harbouring and also apply the appropriate product as a perimeter treatment around the outside of the house if justified. Furniture and soft furnishings may also need to be treated.

The brown dog tick can be difficult to control when active in great numbers. The female is capable of laying up to 4,000 eggs and can survive for up to 18 months without a blood meal. As a result, repeated treatments may be required, and acaricide resistance is a growing issue. This tick species may be able to live on the walls outside the affected house and also outbuildings (especially if dogs spend time in such buildings) so these areas may also require treatment.

Amateur use products from supermarkets and DIY stores etc. are unlikely to be the best route so a professional should always be consulted in cases of brown dog tick activity in domestic properties.

Pest control measures should not be applied in isolation. The dogs within the property must also be treated with appropriate tick treatments, strictly following dosage requirements, and daily tick checking/removal carried out, to ensure that fully fed females cannot lay eggs in the property and start the infestation process again. Infested dogs should not be moved to other dwellings during infestations, to reduce the risk of secondary introductions. Interruption of the life cycle using this multipronged approach will give the greatest chance of success in terms of controlling brown dog ticks in dwellings



Chartered Institute of
Environmental Health



The National Pest Advisory Panel
Chartered Institute of Environmental Health
Chadwick Court, 15 Hatfields, London SE1 8DJ
Telephone 020 7928 6006 Fax 020 7827 5831
Email npap@cieh.org Web www.cieh.org/policy/npap